

Table 1 The inhibition zone diameter (mm) of Bryophytes extract on 11 bacteria (mm)

Test specimen	<i>Staphylococcusepid</i>	<i>Pseudomonas</i>	<i>Vibrio parahemoly</i>	<i>Staphylococcus</i>	<i>Streptococcus</i>	<i>Staphylococcus</i>	<i>Escherichia</i>	<i>Klebsiella</i>	<i>Salmonella</i>	<i>Bacillus</i>	<i>Salmonella</i>
	<i>ermidis</i>	<i>aeruginosa</i>	<i>ticus</i>	<i>aureus</i>	<i>pyogenes</i>	<i>albus</i>	<i>coli</i>	<i>pneumoniae</i>	<i>typhimurium</i>	<i>subtilis</i>	<i>paratyphi-B</i>
<i>Palamocladium euchloron</i>	-	-	-	-	1.50±0.19	-	-	-	-	1.54±0.21	1.46±0.16
<i>Entodon taiwanensis</i>	-	1.54±0.29*	1.56±0.29	1.52±0.29	1.50±0.27	-	1.54±0.30	1.58±0.27	1.57±0.28*	-	1.52±0.26*
<i>Brachythecium moriense</i>	1.53±0.14	1.47±0.13	1.59±0.17	1.54±0.15	-	-	1.59±0.14	1.51±0.14	1.52±0.13	1.56±0.15	1.51±0.15 *
<i>Eurohypnum leptothallum</i>	1.58±0.27	1.52±0.20	-	1.54±0.22	1.58±0.28*	-	-	-	1.56±0.25	-	1.54±0.27 *
<i>Trachypus bicolor</i>	-	1.53±0.19*	-	-	1.55±0.19*	-	-	-	1.55±0.13	-	-
<i>Hypnum calcicolum</i>	1.53±0.20	1.58±0.21**	1.52±0.15	-	-	-	-	-	1.51±0.18	-	1.58±0.26 **
<i>H. fertile</i>	1.54±0.15	1.51±0.12	1.56±0.23	1.53±0.16	-	1.62±0.27	-	1.51±0.13	1.54±0.15	-	1.55±0.19 **
Ethanol	1.52±0.03	1.45±0.06	1.50±0.00	1.49±0.04	1.47±0.05	1.57±0.01	1.51±0.01	1.50±0.02	1.49±0.02	1.53±0.06	1.43±0.06

Note: "-" means that the diameter of the bacteriostatic zone is smaller than that of the control group. The diameter of the bacteriostatic zone is ± standard deviation. \* means  $p<0.05$  compared with the control group, which is significant; \*\* means  $p<0.01$  compared with the control group, which is extremely significant